

Admissible Subspaces and the Denseness of the Intersection of the Domains of Semigroup Generators

PAUL R. CHERNOFF*

Department of Mathematics, University of California, Berkeley, California 94720

AND

JEROME A. GOLDSTEIN†

Department of Mathematics, Tulane University, New Orleans, Louisiana 70118

Communicated by Tosio Kato

Received March 30, 1971

Let $T_i = \{T_i(t) : t \geq 0\}$ be a (C_0) semigroup of linear operators on a Banach space X , with infinitesimal generator A_i , $i = 1, 2$. If Y is a dense Banachable subspace of X left invariant by T_1 , then it is shown that in many cases $D(A_1) \cap Y$ is dense in X , where $D(A_1)$ is the domain of A_1 . In particular, if X is either separable or reflexive and if T_1 leaves $D(A_2)$ invariant, then $D(A_1) \cap D(A_2)$ is dense in X .

1. INTRODUCTION

Let $T = \{T(t) : t \geq 0\}$ be a (C_0) semigroup of linear operators on a Banach space X , and let Y be a dense Banachable subspace of X left invariant by T . While in general it is not true that the restriction of T to Y is strongly continuous when Y is given its Banach space structure, we shall show that in many cases this restriction is strongly continuous for $t > 0$ and the domain of the infinitesimal generator of T intersects Y in a dense subspace of X . As a consequence we obtain a sufficient condition for the intersection of the domains of two (C_0) semigroup generators to be dense.

* Research partially supported by National Science Foundation Grant GP-25082.

† Research partially supported by National Science Foundation Grant GP-19824.

2. TYPES OF SUBSPACES

Let X be a Banach space. A subspace Y of X is *Banachable* if there is a norm which makes Y into a Banach space (of course, this is simply a condition on the cardinality of Y as an abstract vector space). $[Y]$ will denote Y equipped with its Banach space structure. We shall call Y a *type I subspace* of X if Y is a dense Banachable subspace and the natural injection from $[Y]$ to X is continuous. A *type II subspace* of X is a type I subspace Y such that $[Y]$ is homeomorphic to X . A *type III subspace* of X is a type I subspace Y such that $[Y]$ is linearly homeomorphic to X . A *type IV subspace* of X is the domain of a closed densely defined linear operator on X having nonempty resolvent set.

Remarks. 1. A type J subspace is a type $J-1$ subspace, $J = \text{II}, \text{III}, \text{IV}$. This is immediate for $J = \text{II}$ or III . For $J = \text{IV}$, let B be a closed operator having Y as its domain and 0 in its resolvent set. Y is a Banach space under the graph norm of B , and $B : [Y] \rightarrow X$ is the required linear homeomorphism.

2. A subspace Y is type III if and only if it is type IV. We have just seen that type IV subspaces are type III. Conversely, if $Y \subset X$ is type III, let $B : [Y] \rightarrow X$ be a linear homeomorphism. We regard B as an operator on X with domain Y . Then $B^{-1} : X \rightarrow Y$ is bounded. Therefore B is closed and 0 is in its resolvent set.

3. A type II subspace need not be a type III subspace. For example let $X = L^2(0, 1)$, $Y = L^p(0, 1)$, $2 < p < \infty$. Obviously the injection from $[Y]$ to X is continuous, and $[Y]$ is homeomorphic to X (indeed, any two separable Banach spaces are homeomorphic), but Y is not a type III subspace of X .

4. A type I subspace need not be a type II subspace. For example, let $X = L^2(0, 1)$ and $Y = L^\infty(0, 1)$. The point is that here X is separable but $[Y]$ is not. A related example is used in Proposition 1 below. (On the other hand, if X is separable and Y is a type I subspace with $[Y]$ reflexive, it is not difficult to see that $[Y]$ is separable and therefore type II.¹)

¹ Here is a proof. Let $i : [Y] \rightarrow X$ be the inclusion map, let $\{x_n\}$ be a dense sequence in the unit sphere of X , and choose $\varphi_n \in X^*$ with $\|\varphi_n\| = \varphi_n(x_n) = 1$. Then $\{\varphi_n\}$ separates points in X , thus $\{\varphi_n \circ i\} \subset [Y]^*$ separates points in Y . Hence the norm-closed linear span S of $\{\varphi_n \circ i\}$ is a separable weak* dense subspace of $[Y]^*$; consequently $S = [Y]^*$ since $[Y]$ is reflexive. Thus $[Y]^*$ is separable and so is $[Y]$.

3. ADMISSIBLE SUBSPACES FOR SEMIGROUPS

Let $T = \{T(t) : t \geq 0\}$ be a semigroup of bounded operators on a Banach space X which is strongly continuous for $t > 0$, i.e., for each $x \in X$ the map $t \rightarrow T(t)x$ is continuous on $(0, \infty)$. In the terminology of Hille-Phillips [3], T is a (C_0) semigroup if in addition $T(t)x \rightarrow x$ as $t \rightarrow 0$ for each $x \in X$. Whether or not T is (C_0) , the *infinitesimal operator* A_0 of T is defined by

$$A_0x = \lim_{t \rightarrow 0} t^{-1}[T_t x - x]$$

on the domain for which the limit exists. In case T is a (C_0) semigroup, A_0 is closed and densely defined, and is called the *infinitesimal generator* of T . (We shall not make use of the notion of infinitesimal generator for more general classes of semigroups; cf. [3, p. 344].)

According to [3, p. 307] the domain of A_0 and the union of the ranges of the operators $T(t)$ for $t > 0$ have the same closure in X . We shall say that T is an *essential* semigroup provided this closure is the whole of X . Thus any (C_0) semigroup is essential.

Now let T be an essential semigroup on X and let Y be a type I subspace of X . We say that Y is *weakly T -admissible* if $T(t)Y \subset Y$ for each $t > 0$ and if the restriction of T to $[Y]$ is strongly continuous for $t > 0$ and essential. We shall say that Y is *T -admissible* provided $T|_{[Y]}$ is a (C_0) semigroup on Y . We make analogous definitions for one-parameter *groups* of operators; of course for groups there is no difference between admissibility and weak admissibility. The notion of admissibility is useful in several contexts; cf. [2, 4].

PROPOSITION 1. *There exists a Banach space X , a type I subspace Y of X , and a strongly continuous one-parameter group $T = \{T(t) : -\infty < t < \infty\}$ on X which leaves Y invariant, but for which Y is not (weakly) admissible.*

Proof. For $1 \leq p < \infty$ let $X = L_p(-\infty, \infty)$. Define $T(t)f(x) = f(x+t)$ for $f \in X$. Then T is a strongly continuous one-parameter group. Let $Y = \{f \in X : f \text{ is bounded and continuous}\}$. The norm on Y defined by

$$\|f\| = \|f\|_p + \|f\|_\infty$$

makes Y into a type I subspace of X which is obviously left invariant by each $T(t)$. Moreover each $T(t)$ is an isometry on $[Y]$. To see that Y is not T -admissible, let f be any function in Y which is not uniformly continuous on $(-\infty, \infty)$. Then for some $\epsilon > 0$ and all $t \neq 0$ we

have $\|T(t)f - f\| \geq \|T(t)f - f\|_\infty \geq \epsilon > 0$. So strong continuity fails for T on $[Y]$, and hence weak admissibility fails for T on $[Y]$. \square

The next result shows that the distinction between admissibility and weak admissibility is a real one even for (C_0) semigroups T on X .

PROPOSITION 2. *There exists a separable Hilbert space X , a type III subspace Y of X , and a (C_0) semigroup T on X for which Y is weakly admissible but not admissible.*

Proof. We can adapt for this purpose a construction of Hille-Phillips [3, p. 371]. Let X be the space of square-summable sequence pairs $\{(x_n, y_n)\}$ with $\| \{(x_n, y_n)\} \|^2 = \sum_{n=1}^{\infty} (|x_n|^2 + |y_n|^2)$. Let Y be the subspace consisting of all $\{(x_n, y_n)\}$ such that

$$\sum_{n=1}^{\infty} (|x_n|^2 + n|y_n|^2) < \infty.$$

Then Y is a type III subspace of X with $[Y]$ a Hilbert space.

For $t \geq 0$ define $T(t)\{(x_n, y_n)\} = \{(x'_n, y'_n)\}$, where

$$x'_n = e^{-nt}(x_n \cos nt - y_n \sin nt),$$

$$y'_n = e^{-nt}(x_n \sin nt + y_n \cos nt).$$

It is clear that T is a (C_0) contraction semigroup on X . Moreover it is not hard to see that $T(t)Y \subset Y$ for $t > 0$ and that on $[Y]$

$$\|T(t)\| \leq \sup_{n \geq 1} [n^{1/2} e^{-nt}] \leq (2et)^{-1/2}.$$

Thus, $T(t)$ defines a bounded operator on $[Y]$, indeed uniformly bounded for $t \geq \delta > 0$. It is very easy to check that $T|_{[Y]}$ is strongly continuous for $t > 0$. Moreover, the domain of its infinitesimal operator contains all sequences with a finite number of nonzero terms, and is therefore dense in $[Y]$, proving that $T|_{[Y]}$ is an essential semigroup.

However, if $v_k = \{(\delta_{nk}, 0)\}$ and we take $t_k = \pi/2k$ so that $\sin(kt_k) = 1$, $\cos(kt_k) = 0$, we have $T(t_k)v_k = \{(0, \delta_{nk})\}$, whence

$$\|T(t_k)v_k\| = e^{-\pi/2} \sqrt{k} \|v_k\|.$$

This shows that $\|T(t)\|$ is unbounded as $t \rightarrow 0$, so $T|_{[Y]}$ is not of class (C_0) . \square

4. MAIN RESULTS

We shall show now that examples of the type of Proposition 1 are in a sense "pathological," and that under very general hypotheses T -invariant subspaces are weakly T -admissible. The main tool we use is the basic theory of Borel spaces, for which a convenient reference is the book of Parthasarathy [5, Chap. 1].

THEOREM 3. *Let T be an essential semigroup on a separable Banach space X , strongly continuous for $t > 0$. Let Y be a type I subspace of X left invariant by T with $[Y]$ separable. Let $Z \subset X$ be the closure in $[Y]$ of $\bigcup \{T(t)Y : t > 0\}$. Then Z is a weakly T -admissible type I subspace of X .*

Proof. Being complete and separable, X and $[Y]$ are standard Borel spaces. The injection $i : [Y] \rightarrow X$ is continuous, and therefore by [5, p. 21] i is a Borel isomorphism onto its range $Y \subset X$. For $y \in Y \subset X$, $t \rightarrow T(t)y$ is continuous and therefore Borel measurable from $(0, \infty)$ to $Y \subset X$, whence $i^{-1}(T(t)y)$ is Borel measurable from $(0, \infty)$ to $[Y]$. Also each $T(t)$ is a bounded operator on $[Y]$ by the closed graph theorem. Therefore, $T| [Y]$ is a strongly measurable semigroup on Z as defined above, and the infinitesimal operator of $T| [Y]$ has domain dense in Z . So $T| [Z]$ is an essential semigroup strongly continuous for $t > 0$ (cf. [3, pp. 305–307]).

It remains to show that Z is dense in X . Given $x \in X$ we can find a sequence $t_n > 0$ and vectors $y_n \in X$ with $T(t_n)y_n \rightarrow x$, because by assumption T is essential. Since each $T(t_n)$ is bounded and Y is dense in X , we can take the above y_n to be in Y . But then $z_n = T(t_n)y_n \in Z$, so $x \in \bar{Z}$. \square

Remarks. 1. For a simple example where Z is a proper subspace of Y , take $X = L^2(0, 1)$ and $Y = C[0, 1]$ with $T(t)f(x) = x^t f(x)$. Here $Z = C_0[0, 1] = \{f \in Y : f(0) = 0\}$.

Another example is given by $X = L^2(0, 1)$,

$$Y = H^1(0, 1) = \{f \in L^2 : f \text{ is absolutely continuous and } f' \in L^2\}$$

with $\|f\|^2 = \|f\|_2^2 + \|f'\|_2^2$. Here we take $T(t)f(x) = e^{-t/x}f(x)$. It is not hard to verify that $T(t)Y \subset Y$ and $T(t)$ is strongly continuous for $t > 0$ on $[Y]$. Here $Z = H_0^1(0, 1) = \{f \in Y : f(0) = 0\}$.

2. In the first of the above examples Y was not reflexive. In the second it is not hard to see that $T(t)| [Y]$ is not uniformly bounded as $t \rightarrow 0$. In fact if neither of these undesirable things happens, and

if T is a (C_0) semigroup on X , we can deduce that $Y = Z$ and that Y is admissible. This follows from the following result.²

LEMMA 4. *Let Y be a reflexive Banach space. Let $T(t)$ be a semigroup on Y , strongly continuous for $t > 0$, and such that $T(t)y = 0$ for all $t > 0$ implies that $y = 0$. (Note that the latter condition holds if $T(t)$ is the restriction to Y of a (C_0) semigroup on a bigger space X .) Assume in addition that $\|T(t)\|$ is uniformly bounded for t near to 0. Then T is a (C_0) semigroup on Y .*

Proof. Consider the adjoint semigroup $T(t)^*$ on Y^* . The norm $\|T(t)^*\| = \|T(t)\|$ is bounded near 0. Also, $\bigcup \{T(t)^*Y^* : t > 0\}$ is dense in Y^* . Otherwise, since Y is reflexive, there would be a nonzero $y \in Y$ with

$$(T(t)^*y^*)(y) = 0 \quad \text{for all } t > 0, \quad y^* \in Y^*;$$

that is, $y^*(T(t)y) = 0$ so that $T(t)y = 0$ for all $t > 0$. But by assumption this implies $y = 0$.

Note that for all $y^* \in Y^*$, $t \rightarrow T(t)^*y^*$ is weakly continuous for $t > 0$. By [3, p. 306] this implies strong continuity for $t > 0$. The uniform boundedness of $\|T(t)^*\|$ near 0 together with the density of the ranges just proved implies readily that $T(t)^*$ is (C_0) . But then a second dualization implies that $\bigcup \{T(t)Y : t > 0\}$ is dense in Y , so that T is (C_0) on Y . \square

PROPOSITION 5. *Let T be an essential semigroup on X , strongly continuous for $t > 0$, with infinitesimal operator A . Let Y be a weakly T -admissible type I subspace of X . Then $\mathcal{D}^\infty(A) \cap Y$ is dense in X , where $\mathcal{D}^\infty(A) = \bigcap_{n=1} \mathcal{D}(A^n)$ and $\mathcal{D}(A^n)$ is the domain of A^n .*

Proof. Let A_Y denote the infinitesimal operator of the semigroup $T|_Y$. Then $A_Y \subset A$ since the injection from Y to X is continuous. By [3, p. 308] $\mathcal{D}^\infty(A_Y)$ is dense in Y and a fortiori in X . Therefore $\mathcal{D}^\infty(A) \cap Y \supset \mathcal{D}^\infty(A_Y)$ is dense in X . \square

We now come to our main results.

THEOREM 6. *Let X be a Banach space, and let T be an essential semigroup on X , strongly continuous for $t > 0$, with infinitesimal*

² T. Kato has noted that an alternate proof of Lemma 4 can be based on [T. Kato, Remarks on pseudo-resolvents and infinitesimal generators of semi-groups, *Proc. Japan Acad.* **35** (1959), 467–468].

operator A . Let Y be a type I subspace of X left invariant by T . Assume either

(i) X and $[Y]$ are separable

or

(ii) $[Y]$ is reflexive.

Then $\mathcal{D}^\infty(A) \cap Y$ is dense in X .

An immediate consequence of this is the following:

THEOREM 7. Let A_1, A_2 be the infinitesimal generators of (C_0) semigroups T_1, T_2 on a Banach space X which is either separable or reflexive. If T_1 leaves $\mathcal{D}(A_2)$ invariant, then $\mathcal{D}^\infty(A_1) \cap \mathcal{D}(A_2)$ is dense in X .

We simply apply Theorem 6, taking $[Y]$ to be $\mathcal{D}(A_2)$ with the graph norm; the resolvent of A_2 gives an isomorphism from X onto $[Y]$.

The density of $\mathcal{D}(A_1) \cap \mathcal{D}(A_2)$ is a condition that arises in the applications, cf., for instance, DaPrato [1].

Proof of Theorem 6. We first do case (i). By Theorem 3, X has a type I subspace Z ($\subset Y$) which is weakly admissible for T . Hence, by Proposition 5, $\mathcal{D}^\infty(A) \cap Y \supset \mathcal{D}^\infty(A) \cap Z$ is dense in X .

For the proof under assumption (ii) we shall require two lemmas. We will use the following notation: $\|\cdot\|$ is the norm in X , $|\cdot|$ the norm in $[Y]$, B is the unit ball in $[Y]$, and B^* is the unit ball in $[Y]^*$.

LEMMA 8. Let X be a Banach space and Y a type I subspace of X . Let T be a semigroup on X , strongly continuous for $t > 0$, which leaves Y invariant. Then $|T(t)|$ is a locally bounded function on $(0, \infty)$.

Proof. Let $i: [Y] \rightarrow X$ be the natural inclusion map. Then $i^*: X^* \rightarrow [Y]^*$ is a continuous embedding of X^* as a weak* dense subspace of $[Y]^*$. By a well-known result of Banach [3, p. 39], $i^*(X^*) \cap B^*$ is weak* dense in B^* .

We have

$$\begin{aligned} |T(t)| &= \sup\{|T(t)y| : y \in B\} \\ &= \sup\{|\langle \varphi, T(t)y \rangle| : \varphi \in B^*, y \in B\} \end{aligned}$$

and therefore, by the weak* density of $i^*(X^*) \cap B^*$ in B^* ,

$$|T(t)| = \sup\{|\langle i^* \psi, T(t)y \rangle| : y \in B, \psi \in (i^*)^{-1}B^*\}.$$

But for $\psi \in X^*$, $\langle i^*\psi, T(t)y \rangle = \langle \psi, T(t)y \rangle$ is continuous on $(0, \infty)$. Hence $t \rightarrow |T(t)|$, being a supremum of continuous functions, is lower semicontinuous and thus measurable. It is, of course, submultiplicative as well, and therefore bounded on compact subsets of $(0, \infty)$ by [3, p. 241]. \square

LEMMA 9. *Let the hypotheses of Lemma 8 hold, and in addition assume that $[Y]$ is reflexive. Then for each $y \in Y$, the set $\{T(t)y : t > 0\}$ spans a separable subspace of $[Y]$.*

Proof. Given $y \in Y$, let Y_1 be the closed subspace of $[Y]$ spanned by $\{T(t)y : t > 0, t \text{ rational}\}$. By construction Y_1 is separable. We must show that $T(t)y \in Y_1$ for all $t > 0$. Suppose that $T_s y \notin Y_1$ for some $s > 0$. Then there is a $\varphi \in Y_1^*$ such that $\varphi(T(s)y) \neq 0$, but $\varphi(T(t)y) = 0$ for each rational $t \geq 0$. Since $[Y]$ is reflexive, $i^*(X^*)$, which is weak* dense, is actually norm dense in $[Y]^*$. Hence there is a sequence $\{\psi_n\}$ in X^* with $i^*\psi_n \rightarrow \varphi$ in norm in $[Y]^*$. But then

$$\varphi(T(t)y) = \lim_{n \rightarrow \infty} (i^*\psi_n)(T(t)y),$$

and this limit is uniform on compact subsets of $(0, \infty)$ by the local boundedness of $t \rightarrow |T(t)y|$ which follows from Lemma 8. Since the functions $(i^*\psi_n)(T(t)y) = \psi_n(T(t)y)$ are continuous on $(0, \infty)$, the same is true of their uniform limit $\varphi(T(t)y)$. But $\varphi(T(t)y) = 0$ for every rational $t > 0$, hence for every $t > 0$, contradicting $\varphi(T(s)y) \neq 0$. \square

We can now finish the proof of Theorem 6. Suppose that (ii) holds. Let $x \in X$ and $\epsilon > 0$ be given. Since T is essential on X we can find $u \in X$ and $t_0 > 0$ with $\|x - T(t_0)u\| < \epsilon/3$. Since Y is dense in X and $T(t_0)$ is bounded we can find $y \in Y$ with $\|T(t_0)u - T(t_0)y\| < \epsilon/3$. By Lemma 9 the closure in $[Y]$ of the span of $\{T(t)y : t > 0\}$ is separable. Call this space Y_1 , and let X_1 be the closure of Y_1 in the norm of X . Note that X_1 is a closed invariant subspace for T , and that T is essential on X_1 by construction. Hence by case (i) of our theorem, applied to X_1 and Y_1 , there is a $z \in \mathcal{D}^\infty(A) \cap Y_1 \subset \mathcal{D}^\infty(A) \cap Y$ with $\|z - T(t_0)y\| < \epsilon/3$. Hence $\|z - x\| < \epsilon$ and the proof is complete. \square

A final remark: Lemma 9 is not true if the hypothesis that $[Y]$ be reflexive is dropped. To see this let X and T be as in Proposition 1, and let Y consist of all essentially bounded functions $f \in X$ with $|f| = \|f\|_p + \|f\|_\infty$. Let $y \in Y$ be the characteristic function of the interval $[0, 1]$. Then any finite linear combination of rational translates

of y will be a step function whose intervals of constancy all have rational end points. The L^∞ distance from any such function to an irrational translate of y will be at least $1/2$. Hence no irrational translate of y belongs to Y_1 , the closure of the span of $\{T(t)y : t \geq 0, t \text{ rational}\}$.

REFERENCES

1. G. DA PRATO, Somme de générateurs infinitésimaux de class C_0 , *Atti Acad. Naz. Lincei. Rend. Cl. Sci. Fis. Mat. Natur.* **45** (1968), 14–21.
2. J. A. GOLDSTEIN, A Lie product formula for one parameter groups of isometries on Banach spaces, *Math. Ann.* **186** (1970), 299–306.
3. E. HILLE AND R. S. PHILLIPS, "Functional Analysis and Semigroups," American Mathematical Society, Providence, R. I., 1957.
4. T. KATO, Linear evolution equations of hyperbolic type, *J. Fac. Sci. Univ. Tokyo Sect. I* **17** (1970), 241–258.
5. K. R. PARTHASARATHY, "Probability Measures on Metric Spaces," Academic Press, New York, 1967.